



IMPACT OF LEAF CHEMICAL QUALITY INDICES OF DIFFERENT TREE SPECIES ON SOIL MICROBIAL BIOMASS AND N-AVAILABILITY IN A WHEAT MICROCOSM

RAJANI SRIVASTAVA *

Department of Botany, Banaras Hindu University VARANASI - 221005 (U. P.) INDIA

ABSTRACT

Leaves of tropical multipurpose tree species (5 N-fixing, 5 non-N-fixing and combinations of 5 N-fixing species with a non-N-fixing) were incorporated in soil to evaluate effects on soil microbial biomass and N-availability of wheat in a pot experiment under dryland conditions. Soil microbial biomass and N-availability are closely linked to the productivity of an ecosystem. High quality leaves of N-fixing tree species had higher N content, and lower polyphenol (PPL) and lignin (LIG) contents (resulting in lower LIG/N, PPL/N and LIG + PPL/N ratios) than low quality leaves of non-N-fixing species. Combination treatments showed intermediate values of different parameters. Application of high quality leaves caused maximum rise in microbial biomass C and N and available-N in soil; the increase was minimum with low quality leaves and intermediate with combined treatment. Among N-fixing species, *Dalbergia sissoo*, *Cassia fistula* and *Prosopis cineraria* leaf applications showed greater increase in microbial biomass C (128-147% over control) and N (174 - 228%). Strong correlations between LIG + PPL/N ratio of leaves with MBC and MBN of wheat suggests that the ratio can be used to screen multipurpose tree species for use as soil amendment.

Key words: Microbial biomass C and N, *Cassia fistula*, *Dalbergia sissoo*, High quality resource, Low quality resource, *Prosopis cineraria*.

INTRODUCTION

The search for self-sustaining, low-input and energy efficient agricultural system is now a major concern of the researchers and policy makers especially in tropical dryland tracts which account for ca. 68% arable land in India. Sustainable agriculture must focus on biological soil fertility management, relying on a careful synchronization of crop nutrient needs with the availability of those nutrients in the soil^{1,2}. The declining trends of

* Author for correspondence; E-mail: srivastava_252003@yahoo.com: Tel: 09415985202

organic matter content in tropical cultivated soils have necessitated to look for other alternatives such as the application of multipurpose tree leaves, which may serve a dual role as a fertilizer and a source of organic matter. The importance of leguminous trees for soil management has been emphasized in context of application of green leaves to soil as fertilizer³. The suitability of leaf materials as a source of N, may depend to a great extent on its decomposition rate, accumulation of N in soil microbial biomass, and N-mineralization rate in relation to the crop demand. Microbial biomass, a small but active fraction of soil organic matter (1-5% by weight), serves as a reservoir of plant available nutrients⁴ and generally microbial biomass is closely linked to the primary productivity of an ecosystem⁵. Though many studies on leaf/litter decomposition and nutrient release of tropical agroforestry/multipurpose tree species are available^{6,7}, there is lack of studies evaluating the impact of tree leaves on soil fertility indicators. There is a need to screen tropical multipurpose tree species whose leaves can be used as a source of soil amendment; both singly and in combination, particularly in dryland cultivated regions.

The present study aims to evaluate the effect of addition of high and low resource quality leaves on key soil fertility indicators; Soil microbial biomass and available-N of wheat in a pot experiment.

EXPERIMENTAL

Materials and methods

The study was carried out in the Botanical Garden of Department of Botany, BHU (25° 18'N and 83°1' E, 76 m, above sea level). The soil of the study site belongs to the order Inceptisols, sub-order orchrepts, sub-group udic ustocrepts. The fine loamy, mixed, hyperthermic soil neutral to slightly alkaline reaction⁸.

The experiment was designed with leaves of ten tree species used as soil amendments. Amongst the test tree species, 5 were symbiotic N-fixing and 5 non-N-fixing species. The N-fixing species were: *Dalbergia sissoo* Roxb. (Papilionaceae), *Bauhinia variegata* Linn. (Caesalpinaceae), *Cassia fistula* L. (Caesalpinaceae), *Prosopis cineraria* (Linn.) Druce (Mimosaceae) and *Casuarina equisetifolia* J. R. Forst and G. Forst (Casuarinaceae). The non-N-fixing species were: *Sapindus emarginatus* Vahl (Sapindaceae), *Terminalia chebula* Retz. (Combretaceae), *Eucalyptus globulus* Labill. (Myrtaceae), *Madhuca indica* Gmel. (Sapotaceae) and *Holarrhena antidysenterica* (Roth) A. DC (Apocynaceae). Fresh leaves of these species were collected, air dried, and cut into

small (~2 cm) pieces. The substrate in which the test crop, wheat (*Triticum aestivum*, var. HUW 533), was grown comprised field soil. Chopped leaves of 10 above mentioned species and a combination series of five N-fixing species each with non-N-fixing *Terminalia chebula* were incorporated within 0-5 cm deep soil. In all, there were 16 treatments (10 different species + 5 N-fixers with non-N-fixer + control). For each treatment 10 pots were set up. Seeds of the test crop were sown in pots in December 2003 and the crop was harvested after 120 days in April 2004. All pots were placed in an experimental area, which was covered with 3 cm mesh nylon net.

Air-dried tree leaves were finely powdered and the initial chemical composition was determined in triplicate. Carbon and total N content in leaves was determined by ignition method⁹ and microkjeldahl method¹⁰, respectfully. Lignin content (Klason lignin) was estimated¹¹. Extractable polyphenols were determined by Folin-Denis method¹². For the estimation of microbial biomass and available N, soil samples were collected at seedling and maturity stages of crop. Fresh moist soil, subjected to chloroform fumigation-extraction, was used to estimate microbial biomass. Microbial biomass C was measured¹³ and Microbial biomass N was estimated by using microkjeldahl digestion procedure¹⁴. Nitrate-N was measured by the phenol disulphonic acid method¹⁰. Ammonium-N was extracted with 2 M KCl and analysed by the phenate method (APHA 1995)¹⁵.

Treatment mean values were compared using least significant difference (LSD) range test procedure at the 5% level of significance. Correlation between was calculated by using SPSS / PC + software.

RESULTS AND DISCUSSION

Leaves of N-fixing tree species showed lower LIG/N (2.6-10.7), PPL/N (1.0-2.0) and LIG + PPL/N (3.6-12.1) ratios (Table 1). Non-N-fixing species leaves showed distinctly lesser N content but greater lignin and polyphenol contents; therefore, these species showed higher LIG/N (6.3-14.1), PPL/N (1.2-9.2) and LIG + PPL/N (10.4-22.5) ratios. The C/N ratio of N-fixing species leaves (15.8-26.1) was lower than the C/N ratio of non-N-fixing species (23.6-35.7). Combining low N containing *T. chebula* leaves with N-fixing tree leaves provided intermediate values of different parameters. The leaf chemical quality of tree species studied was comparable with characteristics reported in several other multipurpose tree species¹⁶. The leaves of N-fixing species, therefore, represented

potentially rapidly mineralizing high quality resource, and the leaves of non-N-fixing species constituted low quality resource.

Table 1: Leaf quality indices and correlation coefficients (r) of different tree species with soil microbial biomass (MBC, MBN) and available-N; All correlations are significant at $p < 0.01$; for parameter abbreviation see text.

Leaf of tree species	Leaf quality indices			
	C/N	LIG/N	PPL/N	LIG + PPL/N
N-fixing-species				
<i>B. variegata</i>	20.6	8.3	1.9	10.2
<i>C. equisetifolia</i>	26.1	10.7	1.4	12.1
<i>C. fistula</i>	18.6	3.1	1.8	4.6
<i>D. sissoo</i>	15.8	2.6	1.0	3.6
<i>P. cineraria</i>	16.4	4.0	2.0	5.9
Non-N-fixing-species				
<i>E. citriodora</i>	25.2	14.1	5.1	19.2
<i>H. antidysentrica</i>	23.6	5.8	4.6	10.4
<i>M. indica</i>	35.7	13.3	9.2	22.5
<i>S. emarginatus</i>	24.2	9.2	1.2	10.4
<i>T. chebula</i>	25.2	6.3	6.2	12.5
Combined species*				
<i>B. variegata</i> +	22.8	7.4	3.9	11.3
<i>C. equisetifolia</i> +	25.6	8.4	3.9	12.3
<i>C. fistula</i> +	21.4	4.4	3.7	8.1
<i>D. sissoo</i> +	21.5	4.5	3.5	7.9
<i>P. cineraria</i> +	19.9	4.8	3.7	8.5
LSD ($p < 0.05$)	0.47	0.72	0.34	0.77

Cont...

Leaf of tree species	Leaf quality indices			
	C/N	LIG/N	PPL/N	LIG + PPL/N
Soil characteristics				
MBC ($\mu\text{g g}^{-1}$)	-0.73	-0.73	-0.63	-0.80
MBN ($\mu\text{g g}^{-1}$)	-0.81	-0.78	-0.69	-0.86
Available-N	-0.80	-0.73	-0.72	-0.84
*N-fixing species combined with <i>T. chebula</i> ; n = 45				

In all treatments, the soil microbial biomass C and N increased at the maturity stage of wheat (Fig. 1). The increase in microbial biomass towards maturity stage is probably related to the greater nutrient availability in soil due to decreased crop demand and considerable pre-harvest root mortality and decomposition¹⁷. The mean microbial biomass C of five N-fixing species treatments ($291 \mu\text{g g}^{-1}$) was ca.108% higher than the control and the mean of non-N-fixing leaf treatments was 36% greater than control. The mean microbial biomass N showed greater increase (ca. 150% increase relative to control with N-fixing species; ca. 50% increase only with non-N-fixing species). When the non-N-fixing species leaves were combined with low quality *T. chebula* leaves, the mean microbial biomass C and N levels increased 78% and 100%, respectively. Greater increase in the amount of microbial biomass due to N-fixing species leaf application indicates predominant role of N derived from tree leaf decomposition in promoting microbial growth in soil. Substantially greater microbial biomass enhancement with N rich leaves of *D. sissoo*, *C. fistula* and *P. cineraria* (MBC, 128 - 147%; MBN, 174 - 228% over control) supports the above contention. Evidently, microbial biomass N was more strongly correlated with all leaf quality parameters than biomass C (Table 1). Besides, biomass N and available-N exhibited comparable correlations with different leaf quality parameters.

Soil available-N ($\text{NH}_4\text{-N} + \text{NO}_3\text{-N}$) concentrations were greater at crop maturity stage in all treatments (Fig. 1), and $\text{NH}_4\text{-N}$ comprised of 78-82% of available N (data not shown). Greater soil N release occurred in N-fixing species treatments, indicated by 111% greater soil available N concentration than in control (cf. 33% and 67% greater N concentrations in non N-fixing and combined treatments). Addition of *D. sissoo*, *C. fistula* and *P. cineraria* leaves showed >120% increase in soil available N over control, along with greater increase in microbial biomass.




Figure 1. Microbial biomass C, N and available-N ($\mu\text{g g}^{-1}$) at wheat seedling (open bar) and maturity (hatched bar) stages in soil amended with different tree leaves (N-fixing, non-N-fixing and combined species, sequenced as in Table 1); Horizontal lines show mean values for the three groups; vertical lines over the bars show SE of mean; LSD across treatments is shown at $p < 0.05$.

It seems that profuse nutrient release from decomposing leaves of such species not only met the microbial demand but was accumulated in plenty in soil. In non-N-fixing species treatments (excepting *S. emarginatus*) the available N levels remained marginally (ca. 33%) greater than in control. Since incorporation of high quality tree leaves in soil leads to an early net release of N, such leaf resources may serve as major source of nitrogen to meet the immediate requirements during vegetative and reproductive growth phases of quick maturing crops. The addition of poor quality resource results in fierce competition between microbes and crops for available N, with major portion being immobilized by the former; thereby drastically reducing N availability to the crop in the short term¹. Very rapid mineralization has been reported from residues of high quality *Sesbania sesban* and low mineralization from low quality residues of *Grevillea robusta*¹⁸. When mixed with *T. chebula*, *D. sissoo* and *C. fistula* treatments showed ca. 77-80% increase over control, but in case of combined input with *P. cineraria* the increase was much smaller. Combined high + low quality resources (with the three legumes studied presently) showing slower N release can be more suited as N source for slower maturing crops having longer life span, or for the succeeding crops.

There is a need to assess the indicator value of chemical composition of added leaf material accurately to anticipate crop growth and yield enhancements. N concentration and C/N ratio have traditionally been used to assess decomposition and nutrient release potential of organic inputs in natural ecosystems. More recently, lignin and polyphenol contents and their ratios (LIG/N, PPL/N, LIG + PPL/N) have been evaluated for the same purpose¹⁸, though less frequently. Strong correlations found in this study between LIG + PPL/N ratio of incorporated tree leaves with microbial biomass and available N in soil indicate the regulatory effect of interactions among leaf constituents on soil nutrient availability that is directly related to crop growth. Thus, LIG + PPL/N ratio can be used as a tool to screen leaf of potential tree species for crop fertilization.

ACKNOWLEDGEMENTS

We thank Head and the Programme Co-ordinator, Centre of Advanced Study in the Department of Botany for providing laboratory and library facilities. This research was supported by the Pitambar Pant National Environment Fellowship grant from Ministry of Environment and Forests, New Delhi.

REFERENCES

1. R. J. K. Myers, C. A. Palm, E. Cuevan, I. U. N. Gunatilleke, M. Brossard, The Synchronization of Nutrient Mineralization and Plant Nutrient Demand. in, P. L. Wooster, M. J. Swift, (Eds) the Biological Management of Tropical Soil Fertility A Wiley-Sayce Publication, (1994) pp. 81-114.
2. M. Sarrantonio, Soil Response to Surface-Applied Residues of Varying Carbon-Nitrogen Ratios. *Biol Fertil Soils*, **37**, 175-183 (2003).
3. P. W. Chirwa, C. R. Black, C. K. Ong, J. A. Maghembe, Tree and Crop Productivity in Gliricidia/Maize/Pigeonpea Cropping System in Southern Malawi. *Agroforest. Syst.*, **59**, 265-277 (2003).
4. J. L. Smith, E. A. Paul, The Significance of Soil Microbial Biomass Estimations. in, J. M. Bollag, G. Stotzky, (Eds) *Soil Biochemistry Vol. 6*, Marcel Dekker, New York (1990) p. 357-396.
5. H. C. Piao, Y. Y. Wu, Y. T. Hong, Z. Y. Yuan, Soil Released Carbon Dioxide from Microbial Biomass Carbon in the Cultivated Soils of Karst Areas of Southwest China. *Biol Fertil Soils*, **31**, 422-426 (2000).
6. R. Byard, K. C. Lewis, F. Montagini, Leaf Litter Decomposition and Mulch Performance from Mixed and Monospecific Plantations of Native Tree Species in Costa Rica. *Agric Ecosyst Environ.*, **58**, 145-155 (1996).
7. C. A. Palm, R. J. K. Mayers, S. M. Nandwa, Combined Use of Organic and Inorganic Nutrient Sources for Soil Fertility Maintenance and Replenishment. in, R. J. Buresh, P. A. Sanchez, F. Cathoun, (Eds) *Replenishing Soil Fertility in Africa Sssa*. Special Publication No. 51. Soil Science Society of America, American Society of Agronomy, Madison, Wis (1997) pp. 193-217.
8. R. Srivastava, K. P. Singh, Variations in Soil Organic C and N Storage Due to Cultivation Practices in the Gangetic Plain, India. *Int. J. Ecol. Environ. Sci.*, **28**, 193-199 (2002).
9. J. F. Mcbrayer Jr., K. Cromack, Effect of Snow-Pack on Oak-Litter Breakdown and Nutrient Release in A Minnesota Forest. *Pedobiologia*, **20**, 47-54 (1980).
10. M. L. Jackson, *Soil Chemical Analysis*. Prentice-Hall, Inc. Englewood, Cliffs, New Jersey. (1973) p. 485.
11. M. J. Effland, Modified Procedure to Determine Acid Insoluble Lignin in Wood and Pulp. *Technical Association of Pulp Industry (TAPPI)* **60**, 143-144 (1977).
12. J. M. Anderson, J. S. I. Ingram, *Tropical Soil Biology and Fertility, A Handbook of Methods*, Cab International, Wallingford, U.K. (1993)

13. E. D. Vance, P. C. Brookes, D. S. Jenkinson, An Extraction Method for Measuring Soil Microbial Biomass C. *Soil Biol Biochem.*, **19**, 703-707 (1987).
14. P. C. Brookes, A. Landman, G. Pruden, D. S. Jenkinson, Chloroform Fumigation and the Release of Soil Nitrogen, A Rapid Direct Extraction Method to Measure Microbial Biomass Nitrogen in Soil, *Soil Biol. Biochem.*, **17**, 837-842 (1985).
15. Apha, American Public Health Association, Standard Methods for Examination of Water and Wastewater 19th Edition. American Public Health Association, Washington, DC. (1995)
16. R. L. Semwal, R. K. Maikhuri, K. S. Rao, K. K. Sen, K. G. Sexen, Leaf Litter Decomposition and Nutrient Release Patterns of Six Multipurpose Tree Species of Central Himalaya, India. *Biomass and Bioenergy*, **24**, 3-11 (2003).
17. C. P. Kushwaha, S. K. Tripathi, K. P. Singh, Variations in Soil Microbial Biomass and N-Availability Due to Residue and Tillage Management in A Dryland Rice Agroecosystem. *Soil Till Res.*, **56**, 153-166 (2000).
18. G. Nyberg, A. Ekblad, R. Buresh, P. Högberg, Short-Term Patterns of Carbon and Nitrogen Mineralization in A Fallow Field Amended With Green Manures from Agroforestry Trees. *Biol Fertil Soils*, **36**, 18-25 (2002).

Accepted : 22.06.2008