

STUDIES ON ANTI-HIV ACTIVITY AND CYTOTOXICITY OF LEAF OF MORINDA CITRIFOLIA L

PERIYASAMY SELVAM*

Devaki Amma Memorial College of Pharmacy, Chelembra, Pulliparamaba Post, MALAPURAM – 673634 (Kerala) INDIA

ABSTRACT

Different leaf extracts of *Morinda citrifolia* L have been studied for antiviral activity against replication of HIV-1 (IIIB) and HIV-2(ROD) in MT-4 cells. Cytotoxicity of *Morinda citrifolia* against mock-infected MT-4 cells was also assessed by the MTT method. Ether extract (EMC) of *Morinda citrifolia* exhibited a maximum protection of 58 % against replication of HIV-1 in MT-4 cells (EC₅₀=38 μ g/mL and CC₅₀ = 66 μ g/mL). Methanolic extract (MMC) displayed cytostatic activity in MT-4 cells (C-type Adult T cell leukemia cells) with CC50 = 51 μ g/mL.

Key word: Morinda citrifolia, Anti-HIV activity, Cytotoxicity, MTT assay.

INTRODUCTION

Morinda citrifolia L (Noni) is a versatile medicinal plant with a broad spectrum of pharmacological activities. Morinda citrifolia is reported to possess hepatoprotective^{1,2} anticancer³, immunomodulator⁴, anti-inflammatory⁵, wound healing⁶, antioxidant⁷, antitubercular⁸ and wide spectrum of biological activity⁹ and safe herbal drug¹⁰. Recently, much attention has been devoted for searching potential antimicrobial agents from natural products, but anti-HIV activity of morinda citrifolia is relatively less explored. The present work is to study the anti-HIV activity of various extracts of the leaf powder of morinda citrifolia against HIV 1 and 2 in MT-4 cells.

EXPERIMENTAL

Materials and methods

The leaf powder of *morinda citrifolia* are dried under shade and powdered. The powder is extracted with different solvents (acetone, chloroform, methanol, ether and ethyl

^{*}Author for correspondence; E-mail: periyasamy selvam@yahoo.co.in

accetate) for five days by cold maceration. It is then filtered to get the extracts evaporated to dryness under vaccum. The dried extracts acetone (AMC) chloroform (CMC), ether (EMC), ethylacetate (EAMC) and methanol (MMC) were used for anti-HIV activity and cytotoxicty studies.

Anti HIV assay

Morinda citrifolia L. was tested for its inhibitory effects against replication of HIV-1 (IIIB) in MT-4 cells¹¹. The MT-4 cells were grown and maintained in RPMI 1640 DM Medium supplemented with 10% (v/v) heat-inactivated Fetal Calf Serum (FCS), 2 mM-glutamine, 0.1% sodium bicarbonate and 20 μg/mL gentamicin (culture medium). Inhibitory effect of Morinda citrifolia on HIV-1 replications was monitored by inhibition of virus-inducted cytopathic effect in MT-4 cells and were estimated by MTT assay. Briefly, 50 μL of HIV-1 (IIIB) (100-300 CCID50) were added to a flat-bottomed microtiter tray with 50 μL of medium containing various concentrations of Morinda citrifolia. MT-4 Cells were added at a final concentration of 6 x 105 cells/mL. After 5 days of incubation at 37°C, the number of viable cells were determined by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) method. Cytotoxicity of Morinda citrifolia against mockinfected MT-4 cells was also assessed by the MTT method. The anti-HIV activity and cytotxicity data of Morinda citrifolia are presented in Table 1.

RESULTS AND DISCUSSION

Various extract of leaf powder of *Morinda citrifolia*. L has been evaluated for its anti-HIV activity and cytotoxicity (Table 1) against HIV-1 and -2 replication in acutely infected MT-4 cells. *Morinda citrifolia* extracts exhibited a maximum protection of 8-58 against the replication HIV-1 and -2 in acutely infected MT-4 cells. Ether extract of *Morinda citrifolia* (EMC) exhibited antiviral activity with EC₅₀=38 μ g/mL and CC₅₀=66 μ g/mL and 58% maximum protection against the replication HIV-1 in acutely infected MT-4 cells. Methanolic extract of *Morinda citrifolia* (MMC) displayed cytostatic activity against MT-4 cells (Adult T Cell leukemia) with CC₅₀=51.20 ± 15.74 μ g/mL.

The Polynesians utilized the whole Noni plant in various combinations for herbal remedies^{12,13} such as arthritis, diabetes, high blood pressure, muscle aches and pains, menstrual difficulties, headaches, heart disease, AIDS, cancers, gastric ulcers, sprains, mental depression, senility, poor digestion, atherosclerosis, blood vessel problems, and drug addiction. From this study, it has been observed that different extracts of leaf portion of *morinda citrifolia* exhibits cytotoxic activity in MT-4 cells and ether extract (EMC) inhibits

the replication of HIV 1 below the cytotoxic concentration and further studies regarding isolation of active constituents from ether extract is under way.

Table 1: Anti-HIV activity of leaf extracts of morinda citrifolia

Extract	Strain	EC ₅₀ (μg/mL)	CC ₅₀ (µg/mL)	Maximum protection (%)
AMC	IIIB	>138.75	138.75 ± 2.06	10
	ROD	>138.75	138.75 ± 2.06	8
CMC	IIIB	>119.28	119.28 ± 18.35	11
	ROD	>119.28	119.28 ± 18.35	9
EAMC	IIIB	>147	147 ± 19.71	10
	ROD	>147	147 ± 19.71	8
EMC	IIIB	38.50	66 ± 0.06	58
	ROD	>49.70	49.70 ± 14.59	18
MMC	IIIB	>51.20	51.20 ± 15.74	10
	ROD	>51.20	51.20 ± 15.74	15
AZT	IIIB	0.00012	65.90	126
	ROD	0.00062	65.90	148

^a50% Effective concentration of compound, achieving 50% protection of MT-4 cells against the cytopathic effect of HIV.

ACKNOWLEDGMENT

Author thanks Dr. Christophe Pannecouque, Rega Institute for Medical Research, Katholieke Universitiet-Leuven, Belgium for anti-HIV activity.

REFERENCES

- 1. M. Y. Wang, G. Anderson, D. Nowicki and J. Jensen, Hepatic Protection by Noni Fruit Juice Against CCl₄ -Induced Chronic Liver Damage in Female SD Rats, Plant Foods Hum. Nutr., **63(3)**, 141-47 (2008).
- 2. M. Y. Wang, D. Nowicki, G. Anderson, J. Jensen and B. West, Liver Protective Effects of *Morinda Citrifolia* (Noni), Plant Foods Hum. Nutr., **63(2)**, 59-63 (2008).

^b50% Cytotoxic concentration of compound, required to reduce the viability of mock-infected MT-4 cells by 50%

- 3. T. Akihisa, K. Matsumoto, H. Tokuda, K. Yasukawa, K. I. Seino, K. Nakamoto, H. Kuninaga, T. Suzuki and Y. Kimura, Anti-Inflammatory and Potential Cancer Chemopreventive Constituents of the Fruits of *Morinda Citrifolia* (Noni), J. Nat. Prod., **71(7)**, 1322-25 (2008).
- 4. A. K. Palu, A. H. Kim, B. J. West, S. Deng, J. Jensen and L. White, The Effects of *Morinda Citrifolia* L. (Noni) on the Immune System, Its Molecular Mechanisms of Action, J. Ethnopharmacol., **115(3)**, 502-7 (2008).
- 5. A. K. Palu, A. H. Kim, B. J. West, S. Deng, J. Jensen and L. White, Anti-Inflammatory and Potential Cancer Chemopreventive Constituents of the Fruits of *Morinda Citrifolia* (Noni), J. Nat. Prod., **70(5)**, 754-60 (2007).
- 6. A. S. Nayak, G. N. Isitor, A. Maxwell, V. Bhogadi and D. D. Ramdath, Wound-Healing Activity of *Morinda Citrifolia* Fruit Juice on Diabetes-Induced Rats. J. Wound Care, **16(2)**, 83-7 (2007).
- 7. A. N. Su, A. D. Pawlus, H. A. Jung, W. J. Keller, J. L. Mclaughlin and A. D. Kinghorn, Chemical Constituents of the Fruits of *Morinda Citrifolia* (Noni) and their Antioxidant Activity, J. Nat. Prod., **68(4)**, 592-8 (2005).
- 8. J. P. Saludes, M. J. Garson, S. G. Franzblau and A. M. Aguinaldo Antitubercular Constituents from the Hexane Fraction of *Morinda Citrifolia* Linn. (Rubiaceae), Phytother Res., **16**(7), 683-87 (2002).
- 9. A. D. Pawlus and D. A. Kinghorn, Review of the Ethnobotany, Chemistry, Biological Activity and Safety of the Botanical Dietary Supplement *Morinda Citrifolia* (Noni). J Pharm. Pharmacol., **59(12)**, 1587-92 (2007).
- 10. A. J. West, C. J. Jensen and J. Westendorf, Noni Juice is Not Hepatotoxic, World J. Gastroenterol., **12(22)**, 3616-20 (2006).
- 11. A. De Clercq, Rapid and Automated Tetrazolium-Based Colorimeteric Assay for Detection of AntiHIV Compounds, J. Virol. Methods, **20**, 309-321(1988)
- 12. M. Y. Wang, B. J. West, C. J. Jensen, D. Nowicki, C. Su, A. K. Palu and G. Anderson, *Morinda Citrifolia* (Noni), A Literature Review and Recent Advances in Noni Research, Acta Pharmacol. Sin., 23, 1127-41 (2002).
- 13. W. McClatchey, From Polynesian Healers to Health Food Stores, Changing Perspectives of *Morinda Citrifolia* (Rubiaceae), Integr. Cancer Ther., **1**, 110-20 (2002).

Accepted: 12.10.2009