



ISSN (PRINT) : 2320 -1967
ISSN (ONLINE) : 2320 -1975



ORIGINAL ARTICLE

CHEMXPRESS 8(4), 240-249, (2015)

Utilization of some sterols derived from *Jatropha crusa* as a substrate for production of androstenedione (AD) and androstenedione (ADD): maximization using isolated strain of *Absidia corymbifera*

E.M.Ahmed^{1*}, M.I.Mokhtar², A.A.Hamdy¹

¹Chemistry of natural and Microbial Products, National Research Center, Dokki, Cairo, (EGYPT)

²Faculty of pharmacy, Najran university, KSA, (EGYPT)

E-mail: eahmed98@hotmail.com

Abstract : An isolated strain of *Absidia corymbifera* show potentiality to transform some sterols from *Jatropha* to important steroids, Androstenedione (AD) and Androstadienedione (ADD) as a major products. An optimization depends on two levels factorial design as a screening the effective factors from the seven cultural factors was used. The most significant factors found to be the

molasses concentration, time and temperature. These factors were further optimized using three levels factorial design depending on polynomial quadratic equation. © Global Scientific Inc.

Keywords : *Absidia corymbifera*; Androstenedione; Androstadienedione.

INTRODUCTION

Sterols are the natural, organic compounds that are widely distributed in all eukaryotic organisms. The sterols are a part of the cell membranes composition^[21]. Important compounds like steroid hormones and bile acids in humans, brassinosteroids - phytohormones in plants are derived from sterols^[14]. The sterols are involved in important growth and developmental processes in most living organisms^[10]. The most common phytosterols are sitosterol, campesterol and stigmasterol. There are other phy-

tosterols like avenasterol and cycloartenol are relatively present in a smaller amounts^[19].

Androstenedione (AD) is a compound has structural and pharmacological resemblance to testosterone hormone. It can be considered an anabolic steroid. It is common for the anabolic steroids to be used to treat hormone deficiency and muscular atrophy caused by the onset of cancer or HIV^[23]. It is able to increase the level of blood testosterone, the body strength, the lean mass and improve the sexual performance. AD possesses many essential properties of an androgen such as binding to the ligand




Figure 1 : Conversion of sterols to important steroids

binding domain of the androgen receptor, induction of its nuclear translocation, and promotion of myogenic differentiation^[27, 12]. One of the most important use is to act as a starting material for preparation of androgens and anabolic drugs and recently for the production of spironolactone. Other compounds chemically and pharmacologically related to AD.

Androstadienedione (ADD) is used as a precursor for preparing pharmaceutically-interesting steroids such as estradiol or estrone^[2, 25]. It is commercially produced by the microbiological transformation of β -sitosterol and cholesterol^[6].

The microbial cleavage of C-17 side chain of cholesterol and various phytosterols have been reported^[18, 27, 15]. These degradation of C-17 side chain of phytosterols yields AD and ADD; two key intermediates used for commercial production of the majority of medically important steroids. Phytosterols are the abundant source for transformation into these precursors as they are byproducts of many vegetable oil refineries and wood pulp industries^[26]

Production of ADD by biotransformation of sterols has not been reviewed exclusively. However, there is extensive literature on biotransformation of sterols to steroidal compounds^[1, 7, 13, 16, 4, 5].

MATERIALS AND METHODS

Extraction and analysis of the sterols.

The total lipid content of the whole plant parts, leaves, stems, fruits, seeds, was extracted according to the method of Folch et al., 1957. The total plant parts were ground to fine powder. The powder (100gm) was mixed with chloroform/methanol 2:1

to 20 time volume. The whole mixture was homogenized for 2 h on shaking water bath at room temperature. The homogenate was then filtered with funnel and filter paper (whatman no.1). The solvent was washed with distilled water and centrifuged at 4000rpm for 10min. the organic phase was separated. The solvent phase was concentrated using Rota vapor under vacuum. The result was expressed as the percentage of lipids in the composition. Unsaponifiable matter was determined. About 10g of *J. curcas* lipid was placed into a roundbottomed flask and 30mL ethanol and 5mL of aqueous KOH solution were added with some boiling stones into the roundbottomed flask. After refluxing for 1 hour. The heating was stopped and the reaction mixture was transferred into the separating funnel. The flask was washed with 10mL ethanol followed by 20mL warm distilled water and then 20mL cold distilled water. The contents of the separating funnel were left to cool at room temperature, after that 50mL of hexane was added into the separating funnel. After shaking the mixture vigorously for 1min, the mixture was left a few minutes to get two phases. The soap solution phase was converted completely into thesecond separation funnel, and 50mL of hexane was added into the separating funnel. After shaking the mixture vigorously for 1 minute, the mixture was left a few minutes to get two phases. The extractions using 50mL of hexane were repeated five times. The combined extracts in the separating funnel were washed three times with 25mL of 10% (v/v) ethanol, after vigorous mixing the separating funnel; the ethanol layer was drawn off after each wash. The hexane was evaporated to dryness under the vacuum

ORIGINAL ARTICLE

using a rotary evaporator, the drying was completed in a vacuum oven at 75-80°C, and was cooled in a desiccator and was weighed. This residue is considered a substrate for the biotransformation process. The residue was dissolved in 50mL 95% ethanol and finally analyzed for the sterols contents. This extract is considered the substrate. GC analysis was carried out using a Varian 3300 gas chromatography equipped with flame ionization detection system and a non polar capillary column of 60 m × 0.32 mm, film thickness 0.25 µm. The operating conditions were as follow: The temperature was from 260 to 300 °C at 4 °C / min then kept at 300C. Nitrogen gas was used as a carrier with flow rate (1ml/min). the substrate was injected dissolved in CH₂CL₂.

The percent of the common phytosterols, campsterol, β-sitostero, stigmasterol and Δ-avansterol was calculated (TABLE 1).

Isolation of the fungal strains

A soil sample mixed with debris of vegetation (10g) from rural area south of Egypt, was suspended in 90 ml sterile distilled water. The suspension was placed on rotary shaker at 25c for 24 h. thereafter 1 ml of the suspension was placed on plat containing medium composed of (g%): (NH₄)₂SO₄, 0.2; KCl, 0.3; NaCL, 0.1; MgSO₄, 0.2; Kh₂PO₄, 0.02 and 0.1 substrate and chloramphenicol to prevent the bacterial growth. After incubation for 3 days at 25C the growing colonies were purified and transferred to plates of Potato dextrose agar media. The isolates were kept refrigerated. The purified colonies were identified according to the morphological characteristics. The most ptent fungal strain was further identified by molecular techniques.

Screening and the biotransformation process

Fractions of 50 ml of PDA in 250ml conical flasks were inoculated with 2ml of cell suspension

(4x10⁴ cfu) determined by plat count technique. The flasks were then incubated at 25 C for 48h, then the substrate (0.1%) dissolved in 1ml ethanol was added and the incubation conducted to another 24h.

Analysis of the biotransformation product

After completion of the biotransformation time the contents of each flask was extracted with double of its volume of chloroform and checked. The organic phase was separated using separating funnel and was dried over sodium sulfate anhydrous. The solvent phase was evaporated under vacuum to dryness. The residues were considered the test material. The test material were analyzed for the biotransformation products. The biotransformation products were separated by TLC with comparison with authentic under UVlight and after visualization with spraying H₂SO₄ and heating for 10 min where AD took green color and ADD took red color. Biotransformation products were isolated by HPLC, HP, Agilent series 1100 (USA). Shimadzu, PL Hi-Plex pb column). Samples were analyzed under following condition: the flow rate was set at 4.0 ml/min at room temperature, mobile phase was composed of methanol–water in the ratio 90:10 (v:v). The wavelength was set at 238 nm.

Identification and characterization of the fungal strains

The fungal strains were identified according to the morphological characteristics. The identifications relied on colonial and microscopic characteristics^[9, 20, 22, 11].

The medium

Seven different media were examined for production of the desired biotransformation products using the selected fungal species. These medium were given the following codes, MI, MII, MIII, MIV,

TABLE 1 : Assay for the common phytosterols in plant parts of *Jatropha*

The common Phytosterols	Quantity (g/kg)	Percent of sterols in unsap (%)
Campsterols	0.22	3.14
Stigmasterols	0.23	3.21
β-sitosterols	2.68	38.28
Δ ⁵ -avenatserols	0.16	2.18
Total unsap. fraction	7	

MV, MVI and MVII. The media have the following composition (g%): MI (Potato dextrose agar): potato, 2, Dextrose, 2; MII (Dox's medium): Sucrose, 3, NaNO₃, 0.2, K₂HPO₄, 0.1, KCl, 0.05, MgSO₄, 0.05 and FeSO₄, traces; MIII (Yeast extract medium): yeast extract, 3 and peptone, 5; MIV: Molasses, 10, Yeast extract, 0.3, MnSO₄, 0.004, Sod. citrate, 0.001 and K₂HPO₄, 0.4, MV:(Sabroud's dextrose agar): (NH₄)₂SO₄, 0.2 KCl, 0.3, NaCl, 0.1, MgSO₄, 0.2, KH₂PO₄, 0.02 and glucose, 2, MVI (Malt extract medium): Malt extract, 3 and peptone 5, MVII (Corn meal media): Corn infusion, 5.

Maximization

The maximization was carried out through two steps, the first one is screening for seven different factors including the culture conditions. The factors screening determined the most significant factors and it was performed using two levels(+) and (-). To

evaluate seven different cultural parameters 16 runs were applied the number of runs = 2ⁿ⁻³ where n is the number of variables.

In addition, the second step was concerned with prediction of the maximum yields that can be achieved. Box-behnken model was used to optimize the conditions for the most significant factors^[3]. Each factor was evaluated at three levels (+), (0), (-). The relation between the variables was expressed by second order polynomial equation for the data obtained from the runs design. The equation is as follow:

$$Y = b_0 + b_1X_1 + b_2X_2 + b_3X_3 + b_{11}X_1^2 + b_{22}X_2^2 + b_{33}X_3^2 + b_{12}X_1X_2 + b_{13}X_1X_3 + b_{23}X_2X_3$$

Y is the products yields, b₀ is constant, b₁ is the linear coefficient, b₁₁ is the quadratic coefficient, b₁₂ is interaction coefficient and X₁, X₂ are the variables




Figure 2 : Some sterols converting microorganisms and the suitable medium for the conversion process

ORIGINAL ARTICLE

RESULTS AND DISCUSSION

Screening and the microorganism

After screening study for the seven isolated fungal strain, the best one was identified according to molecular techniques as *Absidia corymbifera*. The strain showed the ability to transform about 33% from the substrate to AD and about 12.5% to ADD the other by products were ignored (Figure 2). The genus *Absidia* was described by van Tieghem in 1876. It is one of 4979 general belonging to the

biological kingdom of the Fungi. Members of the genus are saprobics, easily isolated from soil and decaying vegetation. *Absidia* sp is belonged to order Mucorales which is common to perform transformation of foreign material^[24].

Types of transformation medium

Seven growth media of synthetic and natural composition were used to select the most suitable one for formation of AD and ADD as shown in Fi (2 b). The lowest yield was noticed when Dox's medium was used. While, the medium containing mo-




Figure 3 : The optimum time for the conversion of the sterols to AD and ADD




Figure 4 : The ability of the tested microorganism to the substrate concentrations

lasses and some salts was nearly the best. The media containing high concentrations of carbohydrates have negative impact on the biotransformation process, because these types of organic carbon source are preferred by the microorganisms over the sterols. The organic nitrogen compounds influence beneficially the process.

Duration of the biotransformation reaction

The time periods of incubation can affect the reaction outputs. This can be seen when the time of incubation was varied (Figure 3). The optimum time appeared to be at 24 h after substrate addition. Increasing the time after 24 resulted in noticeable decrease for the desired products AD and ADD. The

same effect was notice for incubation periods less than 24 h.

The substrate concentration

The data in Figure (4) indicate that the increasing of the substrate concentration can result in decrease of products formation. This may be attributed to the toxicity of the substrate and the products to the fungal cells which can increase gradually with the gradual rising of the substrate concentration. This effect was will know and one of the inherent problem during the biotransformation processes. These have been reviewed in detail by Malaviya and Fernandes^[24, 7, 17]. The isolation of new microorganisms aims to evaluate robust organisms that can with-

TABLE 2 : Blackett-Burman design and the results for screening of cultural factors

No.	Molasses	Y.E	Sod. citrate	K2HPO4	MnSo4	Tween	Time	ADD	AD
1	40	4	0.01	5	0.05	0.3	24	27.33	18.11
2	40	2	0.01	3	0.05	0.1	24	19.55	16.23
3	10	2	0.01	5	0.03	0.3	24	48.13	21.44
4	10	2	0.01	3	0.05	0.3	24	37.22	22.34
5	10	4	0.01	5	0.05	0.1	24	28.16	19.67
6	10	2	0.01	5	0.05	0.1	12	25.55	16.65
7	40	4	0.01	3	0.05	0.1	12	17.19	14.65
8	10	2	0.01	3	0.03	0.1	12	15.66	12.2
9	10	4	0.01	5	0.03	0.3	12	32.34	18.99
10	40	4	0.01	3	0.03	0.3	24	35.11	16.68
11	40	2	0.01	3	0.03	0.3	12	27.23	15.76
12	10	4	0.01	3	0.05	0.3	12	25.55	16
13	40	4	0.01	5	0.03	0.1	12	19.23	13.16
14	40	2	0.01	5	0.05	0.3	12	24.61	17.89
15	10	4	0.01	3	0.03	0.1	24	32.66	18.87
16	40	2	0.01	5	0.03	0.1	24	19.56	17.88

TABLE 3 : Statistical analysis for the Blackett-Burman design and the results

Source	Sum of Squares		df		Mean Square		F Value		p-value Prob > F	
	ADD	AD	ADD	AD	ADD	AD	ADD	AD	ADD	AD
Model	872.03	95.31	6	7	145.34	13.62	6.17	6.94	0.0082*	0.0069*
A-Molasses	192.24	15.6	1	1	192.24	15.6	8.16	7.96	0.0189*	0.0225*
C-Sod. citrate	0.59	1.13	1	1	0.59	1.13	0.025	0.58	0.8775	0.4688
D-K2HOP4	13.58	6.23	1	1	13.58	6.23	0.58	3.17	0.4672	0.1127
E-MnSO4	38.32	7.65	1	1	38.32	7.65	1.63	3.9	0.2342	0.0838
F-Tween	399.6	2.69	1	1	399.6	2.69	16.96	1.37	0.0026*	0.2752
G-Time	227.71	20.03	1	1	227.71	20.03	9.66	10.21	0.0126*	0.0127*
Residual	212.11	41.99	9	1	23.57	41.99				
Cor Total	1084.14	15.69	15	8		1.96				

ORIGINAL ARTICLE




Figure 5 : The effect of the factors on the conversion process of sterols to ADD

stand their own products.

The maximization steps

Seven different cultural factors were screened for the most effective ones. The factors screening by two levels factorial design namely, (+) and (-) is indicated in TABLE (2). The effect of seven variables were analyzed by multiple regression analysis method and indicated in TABLE (3). The results indicated that, molasses concentration, time and tween are the highly significant factors (<0.05). The effects of the factors on the conversion process are indicated in Figure (5) for AD and in Figure (6) for ADD. Some factors have positive effect and some others have negative effects. The Pareto chart (Fig-




Figure 6 : The effect of the factors on the conversion process of sterols to AD

ure 7) showed a variation between the effects of the factors. This was noticed for each of the produced AD and ADD. According to these results the three factors namely, molasses, time and tween can be chosen for further optimization step.

Box-Behnken model was used to optimize the conditions for the most significant factors. This model is three levels design, each factor was investigated at three levels, (+), (0) and (-1) as shown in TABLE (4). The analysis of variance (ANOVA) was carried out for the results in TABLE (5) for the two products, AD and ADD. The higher values of determination coefficient R^2 0.91, 0.90 for ADD and AD, respectively confirm the effectiveness of the model. The experimental data were fitted to a second order polynomial model obtained by multiple regression

TABLE 4 : The Box-Behnken design and the obtained results for AD and ADD

No.	Molasses (g/l)	Tween (%)	Time (h)	ADD (%)	AD (%)
1	10	0.4	24	73.22	18.33
2	10	0.3	30	68.11	13.34
3	10	0.4	36	57.34	13.87
4	7.5	0.4	30	59.11	14.15
5	5	0.4	36	29.45	9.45
6	7.5	0.5	24	63.56	15.33
7	7.5	0.4	30	47.16	13.22
8	7.5	0.4	30	47.25	13.44
9	7.5	0.3	36	61.67	15.55
10	5	0.3	30	25.44	6.5
11	7.5	0.4	30	62.33	14.99
12	7.5	0.5	36	59.56	12.11
13	5	0.5	30	27.33	6.77
14	7.5	0.3	24	58.44	17.66
15	10	0.5	30	71.55	16.18
16	5	0.4	24	29.45	8.88
17	7.5	0.4	30	65.45	16.45

TABLE 5 : Statistical analysis of Box-Behnken model for AD and ADD

Stat. measure	ADD	AD	Stat measure	ADD	AD
Std. Dev.	7.09	1.37	R-Squared	0.91	0.90
Mean	53.32	13.48	Adj R-Squared	0.81	0.76
C.V. %	13.29	10.15	Pred R-Squared	0.66	0.14
PRESS	1389.75	109.91	Adeq Precision	8.75	9.23




Figure 7 : The Parto chart for some cultural factors affeting conversion of sterols to AD (a) and to ADD (b)

analysis. ANOVA was used to test the adequacy of the model. The high value of R^2 indicates a good consistency between the experimental data and the data from the model. The model equations in coded terms are as follows:

$$AD = +14.45 + 1.27*A + 0.17*B - 3.40*C - 0.11*A*B -$$

$$0.51*A*C + 0.47*B*C - 0.52*A^2 - 0.99*B^2 - 0.55*C^2.$$

$$ADD = +56.20 + 19.82*A + 1.04*B - 2.08*C + 0.39*A*B - 3.97*A*C - 10.81*A^2 + 2.64*B^2 + 1.90*C^2$$

The data that are represented in Figure (8) show the interaction between the three significant factors, molasses concentration, time and temperature. As

ORIGINAL ARTICLE




Figure 8 : The interaction between the most significant factors affection the conversion of sterols to AD and AADD according to Box-Behnken model

shown a maximum value of AD an ADD can be achieved when the terms near 24-30h, molasses 20g/l and temperature 25-30 C.

REFERENCES

- [1] S.Ahmad, S.K.Garg, B.N.Johri; Biotransformation of sterols: Selective and prospects, *Appl.Microbiol.Biot.*, **94**, 1423-1447 (2012).
- [2] H.N.Bhatti, R.A.Khera; Biological transformations of steroidal compounds: *Biochem Microbiol.*, **43**, 1-14.
- [3] G.E.Box, D.W.Behnken; Some new three level designs for the study of quantitative variables, *Technometrics*, **2**, 455-75 (1960).
- [4] M.V.Donova, O.V.Egorova, V.M.Nikolayeva; Steroid 17 production by microorganisms-a review, *Process Biochem.*, **40**, 2253-2262 (2005b).
- [5] M.V.Donova; Transformation of steroids by action bacteria: are view, *Appl.Biochem.Microbiol.*, **43**, 1-14 (2007).
- [6] M.V.Donova, O.V.Egorova; Microbial steroid transformations: current state a review, *Steroids*, **77**, 1267-1290 (2012).
- [7] P.Fernandes, A.Cruz, B.Angelova, H.M.Pinheiro, J.M.S.Cabral; Microbial Conversion of Steroid Compounds: Recent Developments, *Enzyme and microbial technology*, **32**, 688-705 (2003).
- [8] J.Folch, M.Lees, G.H.S.Stanley; A simple method for the isolation and purification of total lipides from animal tissues, *J.Biol.Chem.*, **226**, 497-509 (1957).
- [9] J.C.Gilman; A manual of soil fungi, Oxford and IBH publishing company, Calcutta, (1957).
- [10] M.A.Hartmann; Plant sterols and membrane environment, *Trends in Plant Science*, **3**, 170-175 (1998).
- [11] K.Hoffmann, S.Discher, K.Voigt; Revision of genus *Absidia* (Mucorales Zygomycetes) based on physiological, phylogenetic and morphological characters, thermotolerant *Absidia* spp. form a coherent group, *Mycocladiaceae fam.nov.Mycol Res.*, **111**, 1169-1183 (2007).
- [12] Jasuja, P.Ramaraj, R.P.Mac, A.B.Singh, T.W.Storer, J.Artaza; -4 androstene-3,17-dione [4-androstenedione] binds androgen receptor promotes myogenesis in vitro and Increases serum testosterone levels, fat-free mass, and muscle strength in hypogonadal men. *J.Clin.Endocrinol.Metab.*, **90**(2), 855-863 (2005).
- [13] K.Kieslich; Microbial side-chain degradation of sterols, *J.Basic Microbiol.*, **25**, 461-74 (1985).
- [14] K.Lindsey, M.L.Pullen, J.F.Topping; Importance of plant sterols in pattern formation and hormone signaling, *Trends in Plant Science*, **8**, 1360-1385 (2003).
- [15] Y.Liu, G.Chen, F.Ge, W.Li, L.Zeng, W.Cao; Efficient biotransformation of cholesterol to androsta-1,4-diene-3,17-dione by a newly isolated actinomycete *Gordonia neofelifaecis*, *World J Microbiol.Biotechnol.*, **27**(4), 759-65 (2011).
- [16] S.B.Mahato, S.Garai; Advances in microbial steroid biotransformation, *Steroids*, **332**, 45-62 (1997).
- [17] A.Malaviya, J.Gomes; Androstenedione Production by Biotransformation of Phytosterols, *Bioresource Technology*, **99**, 6725-6737 (2008).
- [18] W.J.Marsheck, S.Kraychy, R.D.Muir; Microbial degradation of sterols, *Appl Microbiol.*, **23**(1), 72-77 (1972)
- [19] K.Matta, A.M.Lampi, J.Petterson, B.M.Fogelfors, V.Piironen, A.Kamal-Eldin; Phytosterol content in seven oat cultivars grown at three locations in Sweden, *Journal of the Science of Food and Agriculture*, **79**, 1021-1027 (1999).
- [20] H.A.Moubasher; Soil fungi in Qatar and other arab countries, Doha university of Qatar, Center for scientific and applied research, 566P (1993).
- [21] V.Piironen, J.Toivo, A.M.Lampi; Plant sterols in cereals and cereal products, *Cereal Chemistry*, **79**, 148-154 (2002).
- [22] S.J.M.Santos, de C.P.Oliveira; Trufem, B F S. Morphological observations on *Absidia corymbifera* and *Absidia blakesleeana* strains preserved under mineral oil, *Mycoses.*, **46**(9-10), 402-6 (2003).
- [23] N.T.Shahidi; A review of the chemistry, Biological action and clinical application anabolic-androgenic steroids, *Clin.Ther.*, **23**(9), 1355-1390 (2001).
- [24] L.L.Smith; Steroids In: Biotechnology (Rehm J H & Reed Geds) Verlag Chemie, Weinhemie, Germany, **6a**, (1984).
- [25] P.Sripalakit, U.Wichai, A.Saraphanchotiwiththaya; Biotransformation of various natural sterols to androstenones by *Mycobacterium* sp. and some steroid-converting microbial strains, *J.Mol.Catal.B:Enzym.*, **41**, 49-54 (2006).
- [26] H.Yang, F.Yan, D.Wu, M.Huo, J.Li, Y.Cao et al.; Recovery of phytosterols from waste residue of soybean oil deodorizer distillate, *Bioresour Technol.*, **101**(5), 1471-6 (2010).
- [27] M.T.Yazdi, F.Malekzadeh, H.Khatami, N.Kamranpour; Cholesterol-degrading bacteria: Isolation, characterization and bioconversion, *World J Microbiol Biotechnol.*, **16**, 103-5 (2000).